



Make Your Silencing Roar  
Ambion® Silencer® Select siRNAs

ASB Applied Biosystems

Learn More >

Science



Magazine

News

Signaling

Careers

Multimedia

Collections

Current Issue

Previous Issues

Science Express

Science Products

My Science

About the Journal

[Home](#) > [Science Magazine](#) > [24 September 1999](#) >

Appenzeller , pp. 2052 - 2053

Science 24 September 1999:  
Vol. 285. no. 5436, pp. 2052 - 2053  
DOI: 10.1126/science.285.5436.2052

## NEWS FOCUS

### PALEONTOLOGY:

## ***T. rex* Was Fierce, Yes, But Feathered, Too**

Tim Appenzeller

### **A new "FEATHERED" fossil, this one a close relative of *Velociraptor*, adds to evidence that many predatory dinosaurs had some kind of plumage**

Dinosaurs live on as sedate green lizards on the Sinclair gas station signs, but for the real beasts, that look was passé long ago. As anyone familiar with *Jurassic Park* can tell you, dinosaurs were brightly colored and awesomely athletic. Now, paleontologists say it's time for yet another makeover, at least for some of *Jurassic Park*'s most menacing characters.

Picture this: Adult *Velociraptors*, savage man-sized hunters with slashing claws, may have been covered in downy feathers, like newly hatched chicks. The same goes for the young of *Tyrannosaurus rex*, and even the full-grown monster may have had a tuft or two.

That's the implication of the stunningly well-preserved fossils unearthed over the last several years from spectacular fossil beds in China: a total of five small dinosaurs apparently clad in some kind of feathers, ranging from fibrous down to true feathers as unmistakable as a pigeon's. The most recent find, which a Chinese team reported last week in *Nature*, is a small, birdlike animal called a dromaeosaur. The 125-million-year-old fossil is an undersized ancestor of *Velociraptor*, which makes it almost certain that the later, larger creatures had similar plumage, says dinosaur paleontologist Mark Norell of the American Museum of Natural History (AMNH) in New York City. "We have as much evidence that *Velociraptors* had feathers as we do that Neandertals had hair," he says.

From the widely scattered perches of this and the other feathered fossils on the dinosaur family tree, paleontologists are realizing that many other dinosaurs on intermediate branches probably also had some kind of feathers, which may have served as insulation. "Feathers are marching their way down" the tree, says Paul Sereno of the University of Chicago--and settling on such unlikely creatures as *T. rex* and its relatives. It's a proliferation that, to some, dramatically underscores the proposed evolutionary link between dinosaurs and birds. But even for ardent proponents of that link, it takes some strenuous mental adjustment to picture such a wide range of dinosaurs in their new clothes.

And a small, energetic band of dissenters is not at all ready to make this kind of adjustment. The halos of fibers found around several of the crucial fossils, including the latest one, are far more likely to be some kind of internal connective-tissue fibers left behind when the flesh decayed than anything related to feathers, according to John Ruben of Oregon State

ADVERTISEMENT

ADVERTISEMENT

Submit your paper today

CLICK HERE

Evolutionary Biology • Bioinformatics and  
Systems Biology • Molecular Biol  
Molecular Biology • Plant Biology  
Development • Biology • Immunol  
**Science Signaling**

University in Corvallis. "We think these are just collagen," he says.

This controversy first erupted when a "feathered dinosaur," *Sinosauropteryx*, emerged 3 years ago from the fossil deposits in China's Liaoning province, where volcanic eruptions some 125 million years ago in the early Cretaceous period entombed a menagerie of ancient animals (*Science*, 1 November 1996, p. 720, and 13 March 1998, p. 1626). The fine-grained sediments had preserved a fibrous mane along the creature's neck and back--"protofeathers" to many paleontologists, internal fibers to some others.

No one could deny the reality of the feathers on two fossils found later in the same deposit, creatures called *Caudipteryx* and *Protoarchaeopteryx* (*Nature*, 25 June 1998, p. 753, and *Science*, 26 June 1998, p. 2051). *Caudipteryx*, the better preserved of the two, had unmistakable feathers on its forelimbs and tail, although it was evidently flightless. But that did not win over the doubters, who think birds originated from reptiles that lived well before the dinosaurs. They agreed that the feathers were real enough but said that based on other features, both creatures were not dinosaurs at all but unusual flightless birds. *Caudipteryx*, says Ruben, is "a Cretaceous turkey, so to speak."

Many paleontologists had a very different view, considering *Caudipteryx* and *Protoarchaeopteryx* as feathered dinosaurs all right, but ones that were somewhat removed from the main line of descent to birds. A better candidate for close cousins of the first birds seemed to be the dromaeosaurs, a group of small predators with a massive claw on each hindlimb and shoulder joints that gave the forelimbs a wide range of motion, a prerequisite for the evolution of flight. But paleontologists had few good fossils of these animals, let alone a view of one with feathers.


Now, Xiao-Chun Wu and his colleagues at the Institute of Vertebrate Paleontology and Paleoanthropology in Beijing have filled that gap with yet another fossil from the Liaoning deposits. The fossil they describe in *Nature*, a 40-centimeter-tall, downy dromaeosaur called *Sinornithosaurus*, "has more bird features than any of the other feathered dinosaurs," says James Clark of George Washington University.

"Everything about the fossil is skewed toward similarity to birds," adds Wu, citing the mobility of the shoulders, the shape of the breastbone, and the proportions of its limbs. Unlike other meat-eating dinosaurs, it has forelimbs nearly as long as its hindlimbs, which may have originally served in "a form of predation associated with grabbing and clutching," says Thomas Holtz of the University of Maryland, College Park, and could easily have evolved into wings. And the downy fringe that appears to cover much of the new dromaeosaur is "a nice confirmation of the existence of feathers--or whatever we want to call them--on dromaeosaurs," notes Holtz.

What *Sinornithosaurus* lacks are complex feathers like those of *Caudipteryx*, which troubles advocates of the bird-dino connection. If *Caudipteryx*--which they view as less birdlike--had true feathers, so should this missing link. This conundrum leads Wu to suggest that the dromaeosaur did have true feathers that somehow weren't preserved, perhaps because the wind blew them away, leaving only the down, before the animal was buried and fossilized. To doubters such as Larry Martin of the University of Kansas, Lawrence, however, the absence of true feathers is telling. "Flight feathers are bigger and have more structure than these fibers," so they should have been preserved along with any down, he argues.

Ruben adds that if this dromaeosaur--supposedly so close to birds--didn't have true feathers, the "down" is probably a chimera as well. He thinks the same is true for *Sinosauropteryx*, the first "feathered" dinosaur, as well as another downy creature, a so-called therizinosaur, reported in the 27 May issue of *Nature*. As Ruben puts it, "All of these things are in all likelihood something like collagen connective fibers."

But if Ruben and his fellow skeptics are wrong about the nature of these fibers, the images of many familiar dinosaurs should be softened with a coating of down. In the family tree of small, meat-eating dinosaurs, *Sinosauropteryx* lies well away from the dromaeosaurs and the putative ancestors of birds. "It's an ordinary ground-running dinosaur," in Holtz's view. So



[To Advertise](#) [Find Products](#)

ADVERTISEMENT

FEATURED JOBS

**AQUATIC BIOGEOCHEMIST**  
University of Montana  
Montana

**Institute for Diabetes,  
Obesity and  
Metabolism/Department of  
Radiology**  
University of Pennsylvania  
Pennsylvania

**Appointment of Executive  
Director**  
National University of  
Singapore  
Singapore

**Director**  
University of Wyoming  
Laramie, WY

**POSTDOCTORAL POSITION**  
University of Pittsburgh  
Cancer Institute  
Pittsburgh, PA

**Research Specialist**  
Johns Hopkins University  
School of Medicine  
Baltimore, MD

**Research Associate**  
Manchester Metropolitan  
University  
Manchester, United  
Kingdom

[More jobs](#)

creatures that perch in between could well have been downy too. Says the AMNH's Norell, "If you're willing to consider the fluffy stuff that's covering the body of *Sinosauropteryx* as feathers, you've got to contemplate *T. rex*, ornithomimids [a group of long-legged, ostrich-sized dinosaurs], and many others as having this kind of plumage at one time in their lives."

Don't start describing *T. rex* to your 4-year-old as a toothy version of Big Bird, though. "Whether an adult *T. rex* had full plumage--well, there's no direct evidence for it, and it might not have been great to have a lot of insulation when you weighed 5 or 6 tons and lived in an environment like Louisiana," says Holtz. "I wouldn't be at all surprised if adult *T. rex* had lost its plumage, although it may have had feathers here and there."

---

**THIS ARTICLE HAS BEEN CITED BY OTHER ARTICLES:**

**Lessons From A Tyrannosaur: The Ambassadorial Role Of Paleontology.**

(2003)

*Palaios* **18**, 475-476

**Phylogenetic Context for the Origin of Feathers.**

S. S. Sumida and C. A. Brochu (2000)

*Integr. Comp. Biol.* **40**, 486-503

[Abstract »](#) [Full Text »](#) [PDF »](#)

*Science*. ISSN 0036-8075 (print), 1095-9203 (online)

[Magazine](#) | [News](#) | [Signaling](#) | [Careers](#) | [Multimedia](#) | [Collections](#) | [Help](#) | [Site Map](#) | [RSS](#)

[Subscribe](#) | [Feedback](#) | [Privacy / Legal](#) | [About Us](#) | [Advertise With Us](#) | [Contact Us](#)

© 1999 American Association for the Advancement of Science. All Rights Reserved.

AAAS is a partner of [HINARI](#), [AGORA](#), [PatientInform](#), [CrossRef](#), and [COUNTER](#).